

The radiative $h^0(A^0) \rightarrow \gamma\gamma$ decays in the two Higgs doublet model.

E. O. Iltan *

Physics Department, Middle East Technical University
Ankara, Turkey

Abstract

We study the radiative $h^0(A^0) \rightarrow \gamma\gamma$ decays in the framework of the general two Higgs doublet model, the so-called model III. We estimate the decay width of the processes at the order of magnitude of $10^{-2} GeV$ for the intermediate values of the Yukawa couplings. We observe that the analysis of the two photon system provide a comprehensive information for the complex nature of the Yukawa couplings and therefore, the source of the CP violation.

*E-mail address: eiltan@heraklit.physics.metu.edu.tr

1 Introduction

The radiative decays $S \rightarrow \gamma\gamma$ (S: scalar, pseudo-scalar particle) are interesting to analyze since the output is the two photon system and it is rich due to two different polarizations of photons, namely the parallel and the perpendicular ones. This leads to two different amplitudes, CP odd and CP even, and the measurement of the photon spin polarization provides comprehensive information about the model parameters existing in the amplitudes. Besides this, such decays are the strong candidates to study the direct or indirect CP violation.

There are various studies on the two photon system in the literature. The decays of mesons into two photons have been analyzed theoretically [1] in various models and experimentally [2] in different collaborations. The decay of the standard model (SM) Higgs boson H^0 into two photons is another candidate for the two photon system as an output and it can provide a valuable information to understand the electroweak symmetry breaking and to predict the model parameters [3, 4]. In the SM, the $H^0 \rightarrow \gamma\gamma$ decay exists at the loop level with the help of all massive particles, which have non-zero electromagnetic charge, running in the loop. The most important contributions come from the top quark and W boson. The branching ratio (BR) has been predicted at the order of the magnitude of 10^{-3} in the framework of SM. The addition of the dimension six and dimension eight fermionic and bosonic operators enhances the SM width of this decay up to one order of magnitude [4].

In the two Higgs doublet model (2HDM) which is the minimal extension of the SM, there appears the well known new Higgs scalars h^0 and A^0 in addition to the SM one, H^0 . The two photon decays of these new particles can provide a considerable information about the model parameters since they occur at least at the one loop level where all the charged fermions run in the loop. On the other hand it is possible to study the CP violating effects since the output is the two photon system and it has two different states, CP odd and CP even.

In the present work we analyze the two photon decays of new Higgs bosons h^0 and A^0 , $h^0(A^0) \rightarrow \gamma\gamma$, in the framework of the model III version of the 2HDM. Since the charged fermions, which are running in the loop, are the source of the processes under consideration there exist various Yukawa couplings which are diagonal in flavor indices and complex in general. In our analysis we take into account only the Yukawa couplings corresponding to the heaviest fermions in each group of particles, up-quarks, down-quarks, charged leptons. We also use the restrictions obtained in the literature for the existing Yukawa couplings and we estimate the decay width of the processes at the order of magnitude of $10^{-2} GeV$ for the intermediate values of the Yukawa couplings. On the other hand, these couplings are complex in general

and a new parameter carrying the imaginary part of the coupling arises. However, the analysis of the CP even $A_{h^0(A^0)}^+$ and CP odd $A_{h^0(A^0)}^-$ amplitudes, which correspond to the parallel and perpendicular spin polarization of photon system, shows that the strength of the complexity of the couplings can be predicted by measuring the relative magnitudes of CP odd and CP even amplitudes. In addition to this the measurement these amplitudes and the BR of the processes provide wide information about the masses m_{h^0} and m_{A^0} .

The paper is organized as follows: In Section II, we present the theoretical expression for the decay widths of $h^0 \rightarrow \gamma\gamma$ and $A^0 \rightarrow \gamma\gamma$ decays, in the framework of the model III. Section 3 is devoted to discussion and our conclusions.

2 The radiative $h^0(A^0) \rightarrow \gamma\gamma$ decays in the two Higgs doublet model

The Higgs bosons h^0 and A^0 can decay into two photon system with the help of the fermion loop, which includes all possible quarks and down leptons. The interaction between the scalar bosons and the fermions are derived by the Yukawa lagrangian and the most general one in the 2HDM reads

$$\begin{aligned} \mathcal{L}_Y = & \eta_{ij}^U \bar{Q}_{iL} \tilde{\phi}_1 U_{jR} + \eta_{ij}^D \bar{Q}_{iL} \phi_1 D_{jR} + \xi_{ij}^{U\dagger} \bar{Q}_{iL} \tilde{\phi}_2 U_{jR} + \xi_{ij}^D \bar{Q}_{iL} \phi_2 D_{jR} \\ & + \eta_{ij}^E \bar{l}_{iL} \phi_1 E_{jR} + \xi_{ij}^E \bar{l}_{iL} \phi_2 E_{jR} + h.c. \quad , \end{aligned} \quad (1)$$

where L and R denote chiral projections $L(R) = 1/2(1 \mp \gamma_5)$, ϕ_i for $i = 1, 2$, are the two scalar doublets, $\bar{Q}_{iL}$ are left handed quark doublets, $U_{jR}(D_{jR})$ are right handed up (down) quark singlets, $l_{iL}(E_{jR})$ are lepton doublets (singlets), with family indices i, j . Here the Yukawa matrices $\xi_{ij}^{U,D}$ and ξ_{ij}^E have in general complex entries. The possible mixing between neutral Higgs bosons H^0 and h^0 is switched off by choosing the parametrization for ϕ_1 and ϕ_2 as

$$\phi_1 = \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 0 \\ v + H^0 \end{pmatrix} + \begin{pmatrix} \sqrt{2}\chi^+ \\ i\chi^0 \end{pmatrix} \right] ; \phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}H^+ \\ H_1 + iH_2 \end{pmatrix} , \quad (2)$$

where only ϕ_1 has a vacuum expectation value;

$$\langle \phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} ; \langle \phi_2 \rangle = 0 , \quad (3)$$

and by considering the gauge and CP invariant Higgs potential which spontaneously breaks $SU(2) \times U(1)$ down to $U(1)$ as

$$V(\phi_1, \phi_2) = c_1(\phi_1^\dagger \phi_1 - v^2/2)^2 + c_2(\phi_2^\dagger \phi_2)^2$$

$$\begin{aligned}
& + c_3[(\phi_1^\dagger \phi_1 - v^2/2) + \phi_2^\dagger \phi_2]^2 + c_4[(\phi_1^\dagger \phi_1)(\phi_2^\dagger \phi_2) - (\phi_1^\dagger \phi_2)(\phi_2^\dagger \phi_1)] \\
& + c_5[Re(\phi_1^\dagger \phi_2)]^2 + c_6[Im(\phi_1^\dagger \phi_2)]^2 + c_7,
\end{aligned} \tag{4}$$

with constants c_i , $i = 1, \dots, 7$. This is the case where the SM (new) particles are collected in the first (second) doublet. Notice that, in the following we will replace $\xi_{ij}^{U,D,E}$ by $\xi_{N,ij}^{U,D,E}$ to emphasize that the couplings are related to the neutral interactions.

Now, we consider the radiative decays of the Higgs bosons h^0 and A^0 , $h^0(A^0) \rightarrow \gamma\gamma$. These processes occur with the internal fermion flow Fig. (1) and the decay amplitudes strongly depend on the Yukawa couplings of the Higgs-fermion interactions. The decay amplitude for these radiative decays can be written in terms of two Lorentz structures:

$$M(S \rightarrow \gamma\gamma) = A_S^+ F_{\mu\nu} F^{\mu\nu} + i A_S^- F_{\mu\nu} \tilde{F}^{\mu\nu}, \tag{5}$$

with $\tilde{F}^{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} F^{\alpha\beta}$ and A_S^+ (A_S^-) are CP even (odd) amplitude:

$$\begin{aligned}
A_S^+ &= \frac{c_1^S}{4\pi m_S^2} \sum_i m_i N_i Q_i^2 g_{ii}^{S+} \{-2 + (-1 + 4x_i^S)(Li_2[f_+] + Li_2[f_-])\}, \\
A_S^- &= \frac{c_1^S}{4\pi m_S^2} \sum_i m_i N_i Q_i^2 g_{ii}^{S-} (Li_2[f_+] + Li_2[f_-]),
\end{aligned} \tag{6}$$

where S is the scalar h^0 or A^0 , $c_1^{h^0(A^0)} = -(I)\sqrt{2}\alpha_e$, $\alpha_e = \frac{1}{137}$, $g_{ii}^{h^0\pm} = (\xi_{N,ii}^F \pm \xi_{N,ii}^{F*})$, $g_{ii}^{A^0\pm} = g_{ii}^{h^0\mp}$, $F = U(D; E)$ for $i = u, c, t$ (d, s, b ; e, μ, τ), N_i is the color factor, $N_i = 3(1)$ for $i = u, c, t, d, s, b$ (e, μ, τ), and Q_i is the charge $Q_i = \frac{2}{3}(\frac{-1}{3}; -1)$ for $i = u, c, t$ (d, s, b ; e, μ, τ). Here the functions $Li_2[z]$ are the PolyLogarithmic function $Li_2[z] = \int_z^0 \frac{\text{Log}[1-t]}{t} dt$ and $f_{\pm} = \frac{2}{1 \pm \sqrt{1-4x_i}}$ with $x_i = \frac{m_i^2}{m_S^2}$. Furthermore, we parametrize the complex Yukawa coupling $\xi_{N,ii}^{U(D,E)}$ as

$$\xi_{N,ii}^{U,D,E} = |\xi_{N,ii}^{U,D,E}| e^{i\theta_{ii}}. \tag{7}$$

Finally the decay width of the processes can be obtained by using the expression

$$\Gamma_S = \frac{m_S^3}{16\pi} (|A_S^+|^2 + |A_S^-|^2) \tag{8}$$

In the decays $h^0 \rightarrow \gamma\gamma$ and $A^0 \rightarrow \gamma\gamma$, the outgoing photons are in one of the possible states given by $F^{\mu\nu} F_{\mu\nu}$ and $F^{\mu\nu} \tilde{F}_{\mu\nu}$ which are even and odd in CP, respectively. These states corresponds to the parallel ($\epsilon_1 \cdot \epsilon_2$) and perpendicular ($\epsilon_1 \times \epsilon_2$) spin polarizations of photons. Eq. (6) shows that the decay amplitude is CP even (odd) for the $h^0 \rightarrow \gamma\gamma$ ($A^0 \rightarrow \gamma\gamma$) decay in the case of real couplings, when the one loop calculations are taken into account. The strength of the complex part in the Yukawa couplings can be estimated by detecting the CP odd (even)

amplitude relative to CP even (odd) one, in other words by studying the ratio $\frac{|A_S^+|^2}{|A_S^-|^2}$, for the $h^0 \rightarrow \gamma\gamma$ ($A^0 \rightarrow \gamma\gamma$) decay. Therefore the precise determination of the photon polarization in the experiments give considerable information about the free parameters of the theory under consideration.

3 Discussion

This section is devoted to the analysis of the decay width and the amplitudes for the CP even and CP odd photon states of the processes $h^0 \rightarrow \gamma\gamma$ and $A^0 \rightarrow \gamma\gamma$. These processes occur at least at the loop order with the help of the internal charged fermions Fig. 1. They are driven by the diagonal Yukawa couplings $\xi_{N,ii}^U$ (i =up quarks), $\xi_{N,ii}^D$ (i =down quarks) and $\xi_{N,ii}^E$ (i =down leptons), which are the free parameters of the model used, and they should be restricted by using the appropriate experimental measurements. At first, we assume that the Yukawa couplings $\xi_{N,ii}^U$, $i = u, c$, $\xi_{N,ii}^D$, $i = d, s$ and $\xi_{N,ii}^E$, $i = e, \mu$, are small compared to $\xi_{N,tt}^U$, $\xi_{N,bb}^D$ and $\xi_{N,\tau\tau}^E$ since the strength of them are related to the masses of fermions denoted by the indices, similar to the Cheng-Sher scenerio [5]. Second, we use the predictions for the couplings $\xi_{N,bb}^D$, $\xi_{N,tt}^U$, obtained by using the constraint region due to the restriction of the Wilson coefficient C_7^{eff} , which is the effective coefficient of the operator $O_7 = \frac{e}{16\pi^2} \bar{s}_\alpha \sigma_{\mu\nu} (m_b R + m_s L) b_\alpha \mathcal{F}^{\mu\nu}$ (see [6] and references therein), in the region $0.257 \leq |C_7^{eff}| \leq 0.439$. Here upper and lower limits were calculated using the CLEO measurement [7]

$$BR(B \rightarrow X_s \gamma) = (3.15 \pm 0.35 \pm 0.32) 10^{-4} , \quad (9)$$

and all possible uncertainties in the calculation of C_7^{eff} [6]. The above restriction ensures to get upper and lower limits for $\xi_{N,bb}^D$, $\xi_{N,tt}^U$ (see [6] for details). At this stage we will make the parametrization

$$\xi_{N,ij}^{U,D,E} = \sqrt{\frac{4G_F}{\sqrt{2}}} \bar{\xi}_{N,ij}^{U,D,E} , \quad (10)$$

where G_F is the fermi constant and use it in the following.

Our numerical calculations are based on the choice intermediate values of the couplings for $C_7^{eff} > 0$, $\bar{\xi}_{N,bb}^D = 30 m_b$, $\bar{\xi}_{N,tt}^U \sim 0.0025 GeV$, respecting the constraints mentioned above. The coupling $\bar{\xi}_{N,\tau\tau}^E$ is the other one which should be restricted and we use the prediction obtained in the analysis of the decay $\tau \rightarrow \mu \bar{\nu}_i \nu_i$, $i = e, \mu, \tau$, in [8]. In this work the BR is estimated at the order of the magnitude of $10^{-6} - 10^{-4}$ for the range of the couplings, $\bar{\xi}_{N,\tau\tau}^E \sim 30 - 100 GeV$ and $\bar{\xi}_{N,\tau\mu}^E \sim 10 - 25 GeV$. In the present work, for the coupling $\bar{\xi}_{N,\tau\tau}^E$, we take the optimum

value $\bar{\xi}_{N,\tau\tau}^E \sim 30 \text{ GeV}$. In addition to this we consider the general case that the couplings are complex (see eq. 7)

In Fig. 2 (3), we present m_{h^0} (m_{A^0}) dependence of the decay width Γ of the $h^0(A^0) \rightarrow \gamma\gamma$ decay, for the intermediate value of the complexity parameter $\sin\theta_{ii} = 0.5$ ($i = t\text{-quark}, b\text{-quark}, \tau\text{-lepton}$) of three Yukawa couplings. The Figure 2 (3) shows that the Γ is at the order of the magnitude of $10^{-3} - 10^{-2} \text{ GeV}$ for the decay $h^0 \rightarrow \gamma\gamma$ ($A^0 \rightarrow \gamma\gamma$) and it is not so much sensitive to the mass of h^0 in the given mass interval (decreases with the increasing values of the mass of A^0). With the possible measurement of this radiative decay a considerable information can be obtained for the mass of the scalar bosons, especially for m_{A^0} .

Fig. 4 (5) is devoted to the $\sin\theta$ dependence of the $\Gamma(h^0(A^0) \rightarrow \gamma\gamma)$ for $m_{h^0} = 85 \text{ GeV}$ ($m_{A^0} = 90 \text{ GeV}$). The solid line represents the case where all the Yukawa couplings are complex and $\sin\theta_{bb} = \sin\theta_{tt} = \sin\theta_{\tau\tau}$, the dashed (small dashed; dotted) line represents the case where $\sin\theta_{tt} = \sin\theta_{\tau\tau} = 0$ ($\sin\theta_{bb} = \sin\theta_{\tau\tau} = 0$; $\sin\theta_{bb} = \sin\theta_{tt} = 0$). We observe that the increasing values of $\sin\theta_{bb}$ ($\sin\theta_{tt}$, $\sin\theta_{\tau\tau}$) causes to enhance (weakly suppress, suppress) the Γ in the $h^0 \rightarrow \gamma\gamma$ decay. For the $A^0 \rightarrow \gamma\gamma$ decay, the increasing values of $\sin\theta_{bb}$ ($\sin\theta_{tt}$, $\sin\theta_{\tau\tau}$) causes to suppress (enhance, suppress) the Γ . In the case that all the Yukawa couplings are complex and have the same value, there is an enhancement (suppression) in the Γ for the increasing values of $\sin\theta$, for the $h^0 \rightarrow \gamma\gamma$ ($A^0 \rightarrow \gamma\gamma$) decay. These results may be effective in the determination of the complexity of the Yukawa couplings.

The ratio of CP even and CP odd amplitudes $R_S = \frac{|A_S^+|^2}{|A_S^-|^2}$ is interesting since it depends strongly on the complex nature of the Yukawa couplings. In Fig. 6 (7), we present m_{h^0} (m_{A^0}) dependence of the ratio R_{h^0} (R_{A^0}) for the decay $h^0(A^0) \rightarrow \gamma\gamma$, for the intermediate value of the parameter $\sin\theta_{ii} = 0.5$. Figure 6 (7) shows that the R_{h^0} (R_{A^0}) is greater than one (less than one) for the given range of mass, for the decay $h^0 \rightarrow \gamma\gamma$ ($A^0 \rightarrow \gamma\gamma$) and increases with the increasing values of the mass h^0 (A^0).

Fig.8 (9) represents the $\sin\theta$ dependence of the ratio R_{h^0} (R_{A^0}) for $m_{h^0} = 85 \text{ GeV}$ ($m_{A^0} = 90 \text{ GeV}$). The solid line represents the case where all the Yukawa couplings are complex and $\sin\theta_{bb} = \sin\theta_{tt} = \sin\theta_{\tau\tau}$, the dashed (small dashed; dotted) line represents the case where $\sin\theta_{tt} = \sin\theta_{\tau\tau} = 0$ ($\sin\theta_{bb} = \sin\theta_{\tau\tau} = 0$; $\sin\theta_{bb} = \sin\theta_{tt} = 0$). We observe that the ratio R_{h^0} (R_{A^0}) is strongly sensitive to the complexity of the Yukawa couplings. For the real couplings $|A_S^-|^2$ vanishes and the ratio R_{h^0} tends to infinity. In such a case the only possible state of two photon system is the parallel polarization. The measurement of the perpendicular polarization for the photon system is a sign for the complexity of the Yukawa couplings. The ratio R_{h^0}

decreases with the increasing values of $\sin\theta$ for all Yukawa couplings. This ratio is larger for the case that only the Yukawa coupling $\xi_{N,tt}^U$ (or $\bar{\xi}_{N,\tau\tau}^E$) is complex. This observation can make it possible to determine the relative strengths of the complexity of different Yukawa couplings. On the other hand, the ratio R_{A^0} vanishes for the real coupling and only the perpendicular polarization state exists for the two photon system. Therefore, for the decay $A^0 \rightarrow \gamma\gamma$, the measurement of the parallel polarization for the photon system is a sign for the complexity of the Yukawa couplings. The ratio R_{A^0} increases with the increasing values of $\sin\theta$ for all Yukawa couplings and it is smaller for the case that only the Yukawa coupling $\xi_{N,tt}^U$ (or $\bar{\xi}_{N,\tau\tau}^E$) is complex. Similar to the $h^0 \rightarrow \gamma\gamma$ decay, this is informative in the determination of the relative strengths of the complexity of different Yukawa couplings.

Now we would like to summarize our results:

- We predict the decay width Γ of the decay $h^0(A^0) \rightarrow \gamma\gamma$ at the order of the magnitude of $10^{-3} - 10^{-2} GeV$ and it is not so much sensitive to the mass of h^0 in the given range (decreases with the increasing values of the mass of A^0) Therefore its measurement can provide valuable information about the $h^0(A^0)$ masses and the Yukawa couplings.
- We study the ratio $R_{h^0(A^0)}$ which is constructed using the CP even $A_{h^0(A^0)}^+$ and CP odd $A_{h^0(A^0)}^-$ amplitudes. For the case of real couplings R_{h^0} (R_{A^0}) tends to infinity (zero) and the only possible state of two photon system is the parallel (perpendicular) polarization. The amount of complexity of the Yukawa couplings can explain the relative magnitude of the perpendicular (parallel) polarization. Therefore the measurement of the photon polarization of the decay $h^0(A^0) \rightarrow \gamma\gamma$ gives a considerable information about the complex nature of the Yukawa couplings and strength of their relative complexity.

Therefore, the experimental analysis of these radiative decays would ensure strong information about the scalar boson masses and the Yukawa couplings existing in the new physics beyond the SM.

4 Acknowledgement

This work has been supported by the Turkish Academy of Sciences in the framework of the Young Scientist Award Program. (EOI-TUBA-GEBIP/2001-1-8)

References

- [1] Wu-jun Huo , Cai-Dian Lu , Zhen-jun Xiao, hep-ph/0302177; ; J.L. Goity, A.M. Bernstein, B.R. Holstein, *Phys. Rev.* **D66**, 076014 (2002); S. Fajfer, P. Singer, J. Zupan, *Phys. Rev.* **D64**, 074008 (2001); R. Akhouri, H. Wang, O. I. Yakovlev, *Phys. Rev.* **D64**, 113008 (2001); A. N. Mitra, *Phys. Lett.* **B473**, 297 (2000); G.G. Devidze, G.R. Dzhibuti, A.G. Liparteliani, *Yad.Fiz.* **59**, 2021 (1996); W. Liu, B. Zhang, H. Zheng, *Phys. Lett.* **B461**, 295 (1999); S. Bar-Shalom, G. Eilam, J. Wudka, A. Soni, *Phys. Rev.* **D59**, 035010 (1999); Chia-Hung V. Chang, Guey-Lin Lin, York-Peng Yao, *Phys. Lett.* **B415**, 395 (1997); I. Caprini, L. Micu, M. Visinescu, *Z. Phys.* **C56**, 225 (1992); M.V. Terentev, *Sov.J.Nucl.Phys.* **16**, 576 (1973); G. L. Lin, J. Liu, Y. P. Yao, *Phys. Rev. Lett.* **64**, 1498 (1990); G. L. Lin, J. Liu, Y. P. Yao, *Phys. Rev.* **D42**, 2314 (1990); S. Herlich, J. Kalinowski, *Nucl. Phys.* **D42**, 2314 (1990); H. Simma, D. Wyler, *Nucl. Phys.* **B344**, 283 (1990); T. M. Aliev, G. Turan, *Phys. Rev.* **D48**, 1176 (1993); G. Hiller, E. Iltan, *Phys. Lett.* **B409**, 425 (1997); G. Hiller, E. Iltan, *Mod. Phys. Lett.* **A12**, 2837 (1997); T. M. Aliev, E. Iltan, *Phys. Rev.* **D58**, 095014 (1998); T. M. Aliev, G. Hiller, E. Iltan, *Nucl. Phys.* **B515**, 321 (1998); M. Boz, E. Iltan, *Phys. Rev.* **D62**, 054010 (2000); .
- [2] CLEO Collaboration, T.E. Coan et al., *Phys. Rev. Lett.* **90**, 101801 (2003); BABAR Collaboration, B. Aubert et al., *Phys. Rev. Lett.* **87**, 241803 (2001); NA31 Collaboration, G.D. Barr et al., *Phys.Lett.* **B351**, 579 (1995); W. Wu, A. Breetwas, D. Green, J. Marraffino, FERMILAB-FN-603 (1993).
- [3] R. Martinez, J.A. Rodriguez, M. Vargas (Colombia, Univ. Nacional). Apr 2000. 8pp. Prepared for 3rd Latin American Symposium on High-Energy Physics (SILAF AE III), Cartagena de Indias, Colombia, 2-8 Apr 2000. e-Print Archive: hep-ph/0004135; J. Novotny, M. Stohr, *Czech.J.Phys.* **49**, 1471 (1999); J.G. Korner, K. Melnikov, O. I. Yakovlev, *Phys.Rev.* **D53**, 3737 (1996); J. M. Hernandez, M. A. Perez, J. J. Toscano, *Phys.Lett.* **B289**, 381 (1992); K. Hagiwara, R. Szalapski, D. Zeppenfeld, *Phys.Lett.* **B318**, 155 (1993); A. De Rujula et.al., *Nucl. Phys.* **B384**, 3 (1992).
- [4] J. M. Hernandez, M. A. Perez, J. J. Toscano, *Phys.Rev.* **D51**, R2044 (1995).
- [5] T. P. Cheng and M. Sher, *Phy. Rev.* **D35**, 3383 (1987).
- [6] T. M.Aliev, E. Iltan, *J. Phys.* **G25** (1999) 989.

- [7] S. Chen, et. al., CLEO Collaboration, *Phys. Rev. Lett.* **87** (2001) 251807.
- [8] E. Iltan H. Sundu, hep-ph/0305005




Figure 1: One loop level diagrams contribute to $\Gamma(h^0(A^0)) \rightarrow \gamma\gamma$ decay in the model III version of 2HDM. Solid lines represent the charged fermions denoted by i , dashed lines represent the h^0 and A^0 fields and curly lines represent the photon fields.




Figure 2: The m_{h^0} dependence of the Γ for the decay $h^0 \rightarrow \gamma\gamma$, for the intermediate value of the complexity parameter $\sin\theta_{ii} = 0.5$ (i=t-quark, b-quark, τ -lepton).




Figure 3: The m_{A^0} dependence of the Γ for the decay $A^0 \rightarrow \gamma\gamma$, for the intermediate value of the complexity parameter $\sin\theta_{ii} = 0.5$ (i=t-quark, b-quark, τ -lepton).




Figure 4: The $\sin\theta$ dependence of the Γ for the decay $h^0 \rightarrow \gamma\gamma$ and $m_{h^0} = 85 \text{ GeV}$. The solid line represents the case where all the Yukawa couplings are complex and $\sin\theta_{bb} = \sin\theta_{tt} = \sin\theta_{\tau\tau}$, the dashed (small dashed; dotted) line represents the case where $\sin\theta_{tt} = \sin\theta_{\tau\tau} = 0$ ($\sin\theta_{bb} = \sin\theta_{\tau\tau} = 0$; $\sin\theta_{bb} = \sin\theta_{tt} = 0$).




Figure 5: The same as Fig. 4 but for the decay $A^0 \rightarrow \gamma\gamma$ and $m_{A^0} = 90 \text{ GeV}$.




Figure 6: The same as Fig. 2 but for the ratio R_{h^0} .




Figure 7: The same as Fig. 3 but for the ratio R_{A^0} .




Figure 8: The same as Fig. 4 but for the ratio R_{h^0} .




Figure 9: The same as Fig. 5 but for the ratio R_{A^0} .